

NuTriVene®

Side Effects/Reactions

Daily Supplement

▶ Restlessness

If restlessness/overactivity occurs in child, reduce Daily Supplement dosage by one-half.

If the Daily Supplement is given too close to bedtime, some children may react to the B vitamins in the supplement by becoming restless. It's generally recommended that you provide the Daily Supplement product two hours prior to bedtime.

Daily Enzyme

▶ Diarrhea

If diarrhea develops, reduce dosage by one-half. If diarrhea continues at reduced-dosage, eliminate Daily Enzyme product.

▶ Spit-Up

If an infant experiences a significant increase in spit-up, reduce dosage by one-half. If this continues at reduced-dosage, eliminate Daily Enzyme product.

▶ Reflux

If an infant or child has been diagnosed with reflux, the Daily Enzyme should not be used as this product may exacerbate this condition.

Note: When these conditions are outgrown by child, Daily Enzyme product can be re-introduced.

NightTime Formula

▶ Sleeplessness

If an infant or child falls asleep and then becomes wide awake after an hour or two, or otherwise experiences a dramatic change in sleep patterns, the NightTime Formula dosage should be reduced by one-half. If child is still not sleeping at reduced-dosage, eliminate NightTime Formula product.

▶ Protein Interaction

While not a health hazard, try to avoid mixing protein sources with this product. Protein sources may interfere with the absorption of nutrients contained in the NightTime Formula.

There are no known drug interactions with Nutrivene-D products.